


CLAIMS

What is claimed is:

1. A compound, including enantiomers, stereoisomers, rotomers and tautomers of said compound, and pharmaceutically acceptable salts, solvates or derivatives thereof, with said compound having the general structure shown in Formula I:



Formula I

wherein:

Z is O, NH or NR¹²;

X is alkylsulfonyl, heterocyclysulfonyl, heterocyclylalkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylcarbonyl, heterocyclylcarbonyl, heterocyclylalkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, alkoxycarbonyl, heterocyclylloxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, alkyaminocarbonyl, heterocyclylaminocarbonyl, arylaminocarbonyl, or heteroarylaminocarbonyl moiety, with the proviso that X may be additionally optionally substituted with R¹² or R¹³;

X¹ is H; C₁-C₄ straight chain alkyl; C₁-C₄ branched alkyl or ; CH₂-aryl (substituted or unsubstituted);

R¹² is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl, heterocyclyl, heterocyclylalkyl, aryl, alkylaryl, arylalkyl, heteroaryl, alkylheteroaryl, or heteroarylalkyl moiety, with the proviso that R¹² may be additionally optionally substituted with R¹³.

R¹³ is hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, carboxy, carbalkoxy, carboxamido, alkoxycarbonylamino, alkoxycarbonyloxy, alkylureido, arylureido, halogen, cyano, or nitro moiety, with the proviso that

the alkyl, alkoxy, and aryl may be additionally optionally substituted with moieties independently selected from R¹³.

P1a, P1b, P2, P3, P4, P5, and P6 are independently:

H; C1-C10 straight or branched chain alkyl; C2-C10 straight or branched chain alkenyl;


C3-C8 cycloalkyl, C3-C8 heterocyclic; (cycloalkyl)alkyl or (heterocyclyl)alkyl, wherein said cycloalkyl is made up of 3 to 8 carbon atoms, and zero to 6 oxygen, nitrogen, sulfur, or phosphorus atoms, and said alkyl is of 1 to 6 carbon atoms;

aryl, heteroaryl, arylalkyl, or heteroarylalkyl, wherein said alkyl is of 1 to 6 carbon atoms;

wherein said alkyl, alkenyl, cycloalkyl, heterocyclyl; (cycloalkyl)alkyl and (heterocyclyl)alkyl moieties may be optionally substituted with R¹³, and further wherein said P1a and P1b may optionally be joined to each other to form a spirocyclic or spiroheterocyclic ring, with said spirocyclic or spiroheterocyclic ring containing zero to six oxygen, nitrogen, sulfur, or phosphorus atoms, and may be additionally optionally substituted with R¹³; and

P1' is H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl, heterocyclyl, heterocyclyl-alkyl, aryl, aryl-alkyl, heteroaryl, or heteroaryl-alkyl; with the proviso that said P1' may be additionally optionally substituted with R¹³.

2. The compound of claim 1, wherein X is selected from the group consisting of:



wherein Alkyl is a C1 to C4 straight or branched chain, and Aryl is a phenyl or substituted phenyl.

3. The compound of claim 2, wherein X is -CO-CH₃.

4. The compound of claim 2, wherein X is -CO-phenyl.

5. The compound of claim 1, wherein P5 and P6 are the same and are:

$-(CH_2)_n-C(O)-R^1$, where $n = 1-4$ and R^1 is OH, O-*t*-Bu, OR^3 , NHR^3 , NH-phenyl or NH-trityl, with R^3 being selected from H, C_1-C_4 straight or branched chain alkyl.

6. The compound of claim 1, wherein P5 and P6 are different and are:

$-(CH_2)_n-C(O)-R^1$, where $n = 1-4$ and R^1 is OH, O-*t*-Bu, OR^3 , NHR^3 , NH-phenyl or NH-trityl, with R^3 being selected from H, C_1-C_4 straight or branched chain alkyl.


7. The compound of claim 5, wherein P5 and P6 are $-CH_2-CH_2-C(O)-O-C(CH_3)_3$ or $-CH_2-CH_2-C(O)-OH$.

8. The compound of claim 6, wherein P5 and P6 are independently selected from $-CH_2-CH_2-C(O)-O-C(CH_3)_3$ or $-CH_2-CH_2-C(O)-OH$.


9. The compound of claim 1, wherein P3 and P4 are the same.

10. The compound of claim 1, wherein P3 and P4 are different.

11. The compound of claim 1, wherein P3 and P4 are independently selected from the group consisting of:




12. The compound of claim 1, wherein P2 is selected from the group consisting of:








wherein n is 0, 1, 2 or 3.

13. The compound of claim 1, wherein P1a and P1b are independently selected from the group consisting of:




14. The compound of claim 1, wherein P1' is selected from the group consisting of:



15. The compound of claim 1, wherein Z is NH and X¹ is H.

16. A compound, including enantiomers, stereoisomers, rotomers and tautomers of said compound, and pharmaceutically acceptable salts or solvates of said compound having the general structure shown in Formula II:



Formula II

wherein:

Z is O, NH or NR¹²;

X is alkylsulfonyl, heterocyclysulfonyl, heterocyclylalkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylcarbonyl, heterocyclylcarbonyl, heterocyclylalkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, alkoxy carbonyl, heterocyclylalkoxy carbonyl, aryloxy carbonyl, heteroaryloxy carbonyl, alkyaminocarbonyl, heterocyclylaminocarbonyl, arylaminocarbonyl, or heteroarylamino carbonyl moiety, with the proviso that X may be additionally optionally substituted with R¹² or R¹³;

R¹² is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl, heterocyclyl, heterocyclylalkyl, aryl, alkylaryl, arylalkyl, heteroaryl, alkylheteroaryl, or heteroarylalkyl moiety, with the proviso that R¹² may be additionally optionally substituted with R¹³;

R¹³ is hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, carboxy, carbalkoxy, carboxamido, alkoxy carbonylamino, alkoxy carbonyloxy, alkylureido, arylureido, halogen, cyano, or nitro moiety, with the proviso that the alkyl, alkoxy, and aryl may be additionally optionally substituted with moieties independently selected from R¹³;

P_{1a}, P_{1b}, P₂, P₃, P₄, P₅, and P₆ are independently:

H; C₁-C₁₀ straight or branched chain alkyl; C₂-C₁₀ straight or branched chain alkenyl;


C₃-C₈ cycloalkyl, C₃-C₈ heterocyclic; (cycloalkyl)alkyl or (heterocyclyl)alkyl, wherein said cycloalkyl is made up of 3 to 8 carbon atoms, and zero to six oxygen, nitrogen, sulfur, or phosphorus atoms, and said alkyl is of 1 to 6 carbon atoms; or

aryl, heteroaryl, arylalkyl, or heteroarylalkyl, wherein said alkyl is of 1 to 6 carbon atoms;

wherein said alkyl, alkenyl, cycloalkyl, heterocyclyl, (cycloalkyl)alkyl and (heterocyclyl)alkyl moieties may be optionally substituted with R¹³ and further


5 wherein said P1 may optionally be a spirocyclic or spiroheterocyclic ring, with said spirocyclic or spiroheterocyclic ring containing zero to six oxygen, nitrogen, sulfur, or phosphorus atoms, and may be additionally optionally substituted with R¹³; and

P1' is H, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl, heterocyclyl, 10 heterocyclyl-alkyl, aryl, aryl-alkyl, heteroaryl, or heteroaryl-alkyl; with the proviso that said P1' may be additionally optionally substituted with R¹³; and




indicates a cyclic ring structure, with the proviso that said cyclic ring structure does not contain a carbonyl group as part of the cyclic ring.

15 17. The compound of Claim 16, wherein said




indicates a five-membered ring.

18. The compound of Claim 16, wherein said



20 indicates a six-membered ring.

19. The compound of claim 16, wherein X is selected from the group consisting of:



wherein Alkyl is a C1 to C4 straight or branched chain, and Aryl is a phenyl or substituted phenyl.

20. The compound of claim 19, wherein X is -CO-CH₃.

5 21. The compound of claim 19, wherein X is -CO-phenyl.

22. The compound of claim 16, wherein P5 and P6 are the same and are: -(CH₂)_n-C(O)-R¹, where n= 1-4 and R¹ is OH, O-*t*-Bu, OR³, NHR³, NH-phenyl or NH-trityl, with R³ being selected from H, C₁-C₄ straight or branched chain alkyl.

10 23. The compound of claim 16, wherein P5 and P6 are different and are: -(CH₂)_n-C(O)-R¹, where n= 1-4 and R¹ is OH, O-*t*-Bu, OR³, NHR³, NH-phenyl or NH-trityl, with R³ being selected from H, C₁-C₄ straight or branched chain alkyl.


24. The compound of claim 22, wherein P5 and P6 are -CH₂-CH₂-C(O)-O-C(CH₃)₃ or -CH₂-CH₂-C(O)-OH.

15 25. The compound of claim 23, wherein P5 and P6 are independently selected from -CH₂-CH₂-C(O)-O-C(CH₃)₃ or -CH₂-CH₂-C(O)-OH.


26. The compound of claim 16, wherein P3 and P4 are the same.

27. The compound of claim 16, wherein P3 and P4 are different.

28. The compound of claim 16, wherein P3 and P4 are independently selected from the group consisting of:



20 29. The compound of claim 16, wherein P2 is selected from the group consisting of:



wherein $n = 0, 1, 2, \text{ or } 3$; and

$R^2 = R^3 = H$; $R^2 = C_1 \text{ to } C_6 \text{ straight chain alkyl or cycloalkyl}$; $R^3 = H$
 $R^4 = COAlkyl \text{ (straight chain or cyclic, } C_1 \text{ to } C_6)$; $COAryl$; $COOAlkyl$; $COOAryl$
 $R^5 = H$; $R^6 = Alkyl \text{ (} C_1 \text{ to } C_3)$; $R^6 = H$; $R^5 = Alkyl \text{ (} C_1 \text{ to } C_3)$
 $R^7 = H$; $R^8 = Alkyl \text{ (} C_1 \text{ to } C_3)$, CH_2OH ; $R^8 = H$; $R^7 = Alkyl \text{ (} C_1 \text{ to } C_3)$, CH_2OH ;

$R^7 = R^8 = \text{Alkyl (C}_1 \text{ to C}_3\text{), CH}_2\text{OH}$

$R^9 = R^{10} = \text{Alkyl (C}_1 \text{ to C}_3\text{); } R^9 = \text{H, } R^{10} = \text{Alkyl (C}_1 \text{ to C}_3\text{), COOMe, COOH, CH}_2\text{OH;}$

$R^{10} = \text{H, } R^9 = \text{Alkyl (C}_1 \text{ to C}_3\text{), COOMe, COOH, CH}_2\text{OH;}$

$R^{11} = \text{Alkyl (C}_1 \text{ to C}_6 \text{ straight chain, branched or cyclic), CH}_2\text{Aryl (may be substituted)}$


$Z^1 = Z^2 = \text{S, O; } Z^1 = \text{S, } Z^2 = \text{O; } Z^1 = \text{O, } Z^2 = \text{S; } Z^1 = \text{CH}_2, Z^2 = \text{O; } Z^1 = \text{O, } Z^2 = \text{CH}_2;$

$Z_1 = \text{S, } Z_2 = \text{CH}_2; Z^1 = \text{CH}_2, Z^2 = \text{S}$


$Z^3 = \text{CH}_2, \text{S, SO}_2, \text{NH, NR}^4$

$Z^4 = Z^5 = \text{S, O}$

30. The compound of claim 16, wherein P1a and P1b are independently selected from the group consisting of:




30. The compound of claim 16, wherein P1' is selected from the group consisting of:





- 32
31. The compound of claim 16, wherein Z is NH.
32. A pharmaceutical composition comprising as an active ingredient a compound of claim 1 or claim 16.
33. The pharmaceutical composition of claim 32 for use in treating disorders associated with Hepatitis C virus.
34. The pharmaceutical composition of claim 32 additionally comprising a pharmaceutically acceptable carrier.
35. A method of treating disorders associated with the HCV protease, said method comprising administering to a patient in need of such treatment a pharmaceutical composition which composition comprises therapeutically effective amounts of a compound of claim 1 or claim 16.
36. The method of claim 35, wherein said administration is via subcutaneous administration.
37. The use of a compound of claim 1 or claim 16 for the manufacture of a medicament to treat disorders associated with the HCV protease.
38. A method of preparing a pharmaceutical composition for treating disorders associated with the HCV protease, said method comprising bringing into intimate

contact a compound of claim 1 or claim 16 and a pharmaceutically acceptable carrier.


39. A compound exhibiting HCV protease inhibitory activity, including enantiomers, stereoisomers, rotamers and tautomers of said compound, and pharmaceutically acceptable salts or solvates of said compound, said compound being selected from the group of compounds with structures listed below:




FOI 29060650






T06T20" 29060660







09909062-071901







069066-0701





09909062 071901






09009062 071901




106170" 29060660

173







00909062.071901






090906Z 071901



176








T06T0" 29060660











T06TQ" 29060660










09909062-071901







FOI 2012-07190







09909062.071901



5

~~40.~~ A pharmaceutical composition for treating disorders associated with the HCV protease, said composition comprising therapeutically effective amount of one or more compounds in claim 39 and a pharmaceutically acceptable carrier.

~~41.~~ The pharmaceutical composition of claim 40, additionally containing an antiviral agent.



~~42.~~ The pharmaceutical composition of claim 40 or claim 41. still additionally containing an interferon.

~~43.~~ The pharmaceutical composition of claim 42, wherein said antiviral agent is ribavirin and said interferon is α -interferon.



~~44.~~ A compound selected from the group consisting of:

FOI b7D b7C b6 b5 b4 b3 b2 b1



5





10





15





20





25





30





5





10





15





20



25





30





T06T70"29060660



5





10





15



20















25





30

FOOTNOTES 29060660





T06T20" 29060660



5





10





15





20



25





30





09909060 071901



5





10




15




20













25
















30






10950607190123060660





and



or an enantiomer, stereoisomer, rotamer or tautomer thereof, or a

pharmaceutically acceptable salt or solvate thereof, wherein the compound exhibits HCV inhibitory activity.

45. A pharmaceutical composition, comprising one or more compounds of claim 44.

46. A method of treatment of an hepatitis C virus associated disorder, comprising administering an effective amount of one or more compounds of claim

44.

48
47. A method of modulating the activity of hepatitis C virus (HCV) protease, comprising contacting HCV protease with one or more compounds of claim 44.



49. A method of treating, preventing, or ameliorating one or more symptoms of hepatitis C, comprising administering an effective amount of one or more compounds of claim 44.

50. The method of claim 47, wherein the HCV protease is the NS3/NS4a protease.

51. The method of claim 49, wherein the compound or compounds inhibit HCV NS3/NS4a protease.

52. A method of modulating the processing of hepatitis C virus (HCV) polypeptide, comprising contacting a composition containing the HCV polypeptide under conditions in which the polypeptide is processed with one or more compounds of claim 44.

53. The compound of claim 17, wherein P2 is selected from the group consisting of:



wherein:

n is 0, 1, 2 or 3;

R²⁰ is alkylene-COOH;

R²¹ is C(O)alkyl, CO₂alkyl, C(O)aryl, CO₂aryl, SO₂alkyl, SO₂aryl, CONHalkyl, or CONHaryl;

R²² is alkyl or alkylene-COOH; and

R²³ is alkyl.

54

~~53.~~

The compound of claim 52, wherein:

 R^{20} is CH_2COOH ; R^{21} is CO_2Ph , COPh , CO_2CH_2 -9-fluorenyl, CO -(3-phenoxyphenyl), SO_2Ph or CONHPh ;

5

 R^{22} is methyl or CH_2COOH ; and R^{23} is methyl.

T06T20"29060660